



Tobacco Use and

United States Students

What is the problem?

The 2007 national Youth Risk Behavior Survey indicates that among U.S. high school students:

Cigarette Use

- 50% ever tried cigarette smoking, even one or two puffs.
- 20% currently smoked cigarettes. (1)
- 8% smoked cigarettes on 20 or more days during the 30 days before the survey.
- 6% smoked cigarettes on school property. (2)
- Among students who currently smoked cigarettes, 50% tried to quit smoking cigarettes during the 12 months before the survey.

Other Tobacco Use

- 8% currently used smokeless tobacco. (3)
- 5% used smokeless tobacco on school property. (2)
- 14% currently smoked cigars, cigarillos, or little cigars. (4)

Any Tobacco Use

- 26% reported current cigarette use, current smokeless tobacco use, or current cigar use.

What are the solutions?

Better health education • More family and community involvement
Healthier school environments • More comprehensive health services

What is the status?

The School Health Policies and Programs Study 2006 indicates that among U.S. high schools:

Health Education

- 69% required students to receive instruction on health topics as part of a specific course.
- 47% taught 17 tobacco-use prevention topics in a required health education course.

Family and Community Involvement

- 37% had a school health council that addressed tobacco-use prevention.
- 33% involved students' families and 48% involved community members in the development, communication, and implementation of policies or activities related to tobacco-use prevention.
- 37% had or participated in a youth empowerment or advocacy program related to tobacco-use prevention.

School Environment

- 66% prohibited all tobacco use in all locations. (5)
- 88% prohibited all tobacco advertising. (6)
- 60% posted signs marking a tobacco-free school zone.

Health Services

- 57% provided tobacco-use prevention services at school in one-on-one or small-group sessions.
- 29% provided tobacco-use prevention services to students through arrangements with providers not located on school property.
- 46% provided tobacco-use cessation services at school.
- 27% provided tobacco-use cessation services to students through arrangements with providers not located on school property.

1. Smoked cigarettes on at least 1 day during the 30 days before the survey.
2. On at least 1 day during the 30 days before the survey.
3. Used chewing tobacco, snuff, or dip on at least 1 day during the 30 days before the survey.
4. Smoked cigars, cigarillos, or little cigars on at least 1 day during the 30 days before the survey.
5. Prohibited all tobacco use by students, faculty and school staff, and visitors; in school buildings, outside on school grounds (including parking lots and playing fields), on school buses or other vehicles used to transport students, and at off-campus, school-sponsored events.
6. Prohibited tobacco advertising in school buildings, on school grounds, on school buses or other vehicles, in school publications, and through sponsorship of school events, and prohibited students from wearing tobacco brand name apparel or carrying merchandise with tobacco company names, logos, or cartoon characters.

Where can I get more information? Visit www.cdc.gov/healthyyouth or call 800 CDC-INFO (800-232-4636).



DEPARTMENT OF HEALTH AND HUMAN SERVICES
CENTERS FOR DISEASE CONTROL AND PREVENTION

