

Smoking and Smoking Cessation in England 2010: Findings from the Smoking Toolkit Study

Robert West and Jenny Fidler
Cancer Research UK Health Behaviour Research Centre
University College London

Published 2011

(Citation: West R, Fidler J (2011) *Smoking and Smoking Cessation in England 2010*. London: Vasco-Graphics. Available from www.smokinginengland.info)

This report describes key findings from the Smoking Toolkit Study (STS) relevant to development of tobacco control policy for the years 2007 to 2010. The STS is a continuing series of monthly surveys of representative sample of the population of England aged 16+. Each monthly wave is followed up for 6 months by means of a postal survey. Full details of the study methods are described elsewhere¹. This document focuses on top level study findings relating to smokers and recent ex-smokers. It does not stratify findings by sociodemographic variables. Neither does it address issues relating to harm reduction. A number of papers have been published from the study that include analyses relating to such stratification and to harm reduction²⁻¹⁹

Smoking prevalence

Cigarette smoking prevalence declined from 24.1% in 2007 to 21.4% in 2010 (Figure 1). Prevalence of smoking of any tobacco product in 2010 was 21.9%. Prevalence of any smoking within the past year was 23.3%. There has been little change in smoking prevalence since 2008.




Figure 1: Prevalence of smoking. Base: All respondents aged 16+; N=22,079(2007), 18,991(2008), 21,138(2009), 24,795(2010)

Cigarette consumption and spending on tobacco products

The average daily cigarette consumption has declined from 14.5 cigs per day in 2007 to 13.1 in 2010 ($p < 0.001$ by analysis of variance, Figure 2). There was a sharper fall in consumption of manufactured cigarettes and a rise in the consumption of hand-rolled cigarettes ($p < 0.001$ by analysis of variance in both cases).




Figure 2: Mean daily consumption of different types of cigarette. Base: all current cigarette smokers, N=8,650(2007), 6,446(2008), 6,827(2009), 6,376(2010). Samples sizes are reduced for manufactured and hand-rolled cigarettes because of missing values. This also leads to these not adding exactly to the total number of cigarettes smoked.

The average price paid per 20 cigarettes smoked in 2007 increased from £1.86 for hand rolled and £4.82 for manufactured to £2.08 for hand-rolled and £5.24 for manufactured in 2010. The overall price per cigarette also rose from £3.83 to £4.00 ($p < 0.0001$ for the increase in all cases by linear regression, Figure 3). The increase in the overall price per cigarette was reduced by switching from manufactured to hand-rolled (see Figure 2). This, together with the overall drop in consumption, meant that the weekly cost of smoking remained static from 2008 through 2010 (Figure 4).




Figure 3: Amount spent per 20 cigarettes by type of cigarette and year. Base: All cigarette smokers. N=4,043(2008), 2,238(2009), 2,164(2010)2238(2010). Questions about price paid were not asked in all waves and some missing data for hand-rolled and manufactured cigarettes questions..




Figure 4: Mean weekly spending on cigarettes. Base: All current cigarette smokers. N=4,051(2008), 2,251(2009), 2,171(2010). Question about price paid were not asked in all waves.

The consistency of the information provided by respondents on price paid per week on smoking and numbers of manufactured and hand-rolled cigarettes was assessed by examining the variance accounted for (R^2) in each year when the weekly spending was regressed on to the numbers of different types of cigarette smoked, forcing the regression through the origin. The figures were 0.83, 0.86 and 0.83 for 2008, 2009 and 2010 respectively, showing a very high level of consistency between the different types of information.

Motivation to stop smoking

Motivation to quit is strongly predictive of quit attempts in the subsequent 3 and 6 months ($p < 0.0001$ by chi-squared test for both 3- and 6-month follow up, Figure 5).




Figure 5: Association between ratings of motivation to quit at baseline and reports of quit attempts between baseline and 3-month follow up and 6-month follow up. Base: Current cigarette smokers at baseline who were successfully followed up; N=1708(3 month), 2088(6 month)

The proportion of smokers not wanting to quit has increased from 27.2% in 2008 to 33.6% in 2010. The decrease has been mainly in the percentage who really want to quit but do not know when ($p < 0.0001$ by chi-squared test for overall change in percentages, Figure 6).




Figure 6: Percentage of smokers with different degrees of motivation to quit. Base: Current cigarette smokers at baseline. $N=361(2008), 4553(2009), 5293(2010)$. The question was introduced in 2008.

Attempts to stop smoking

The proportion of smokers who tried to quit in the previous 12 months has declined from 42.5% in 2007 to 35.9% in 2010 ($p < 0.0001$ by chi-squared test, Figure 7).




Figure 7: Percentage of smokers who tried to quit in the preceding 12 months. Base: All respondents who reported having smoked in the past 12 months. $N=21,217$.

Receipt of advice to stop smoking

Only a minority of smokers recalled having discussed smoking with their GP in the past 12 months in 2010 and only 25.9% recalled having been offered a prescription or advised to see a stop smoking practitioner (Figure 8).




Figure 8: Percentage of smokers who reported having received advice from their GP on smoking in the past 12 months. Base: 5,801 respondents in 2010 who had smoked in the past 12 months. Data collection began in 2010

Recollection of having been advised to stop smoking by their GP was not associated with a significant increase in reports of having made a quit attempt, but having received an offer of a prescription or recommendation to see a stop smoking advisor (practice nurse or go to the Stop Smoking Service) was associated with a markedly higher proportion of reports of quit attempts ($p < 0.0001$ by chi-squared test, Figure 9).




Figure 9: Percentage of smokers who reported having made a quit attempt in the past 12 months as a function of advice received from their GP during that time. Base: All respondents who had smoked in the past 12 months, N=21,271

Use of aids to cessation

The proportion of smokers who made at least one quit attempt who used some form of cessation aid increased from 49.1% in 2007 to 53.6% in 2010; the increase was primarily in those who used a prescription with minimal behavioural support ($p < 0.001$ by chi-squared test for the overall change in proportions, Figure 10).




Figure 10: Percentage of smokers who made at least one quit attempt in the past year who used different aids to cessation. Base: All smokers who reported having made at least one quit attempt in the past 12 months. $N=2,534(2007)$, $1,831(2008)$, $1,833(2009)$, $2,068(2010)$




Figure 11: The proportion of smokers who made quit attempts in the past 12 months who used different types of medication. Base: All those who smoked in the past 12 months who made at least one quit attempt in the past 12 months. $N=2,533(2007)$, $1,830(2008)$, $1,832(2009)$, $2,068(2010)$

There has been an increase in proportion of smokers who make quit attempts who used varenicline from 0.06% in 2007 to 8% in 2010 ($p < 0.0001$ by chi-squared test) with no commensurate fall in the use of prescriptions for other medications (Figure 11).

Approaches to quitting

There was a decline in the percentage of quit attempts that were unplanned from 58.1% in 2007 to 48.1% in 2010 ($p < 0.0001$ by chi-squared test, Figure 12). The proportion of quit attempts that involved cutting down gradually remained static at around 40%.




Figure 12: Percentage of smokers adopting different approaches to quitting in their most recent quit attempt. Unplanned=made the quit attempt as soon as the decision was taken and did not even wait a day; gradual=made the quit attempt by cutting down first. Base=All those who smoked in the past 12 months who made at least one quit attempt. $N=2,525(2007), 1,827(2008), 1,832(2009), 2,054(2009)$

Success at stopping smoking

Figure 13 shows the proportion of ever-confirmed-smokers (people who report that they ever smoked for at least a year) who go on to become ex-smokers (not smoked for at least a year), aggregating data from all years of the study. Only 25% achieve ex-smoker status by the age of 35, when they begin to lose significant life-expectancy. Only 55% achieve this by the age of 65.




Figure 13: Percentage of ever-confirmed-smokers (currently smoke or have smoked for at least a year) who were ex-smokers (reported not having smoked for at least a year) as a function of age. Base: All ever-confirmed smokers. $N=28,207$. Data only used for respondents up to the age of 70 because of significant bias from death after that age.

The proportion of last-year smokers (people who had smoked within the past 12 months) who reported no longer smoking declined from 6.7% in 2007 to 4.8% in 2010 ($p<0.001$ by chi-squared test, Figure 14). There was also a fall in the proportion who had succeeded given that they made a quit attempt ($p=0.02$ by chi-squared test).




Figure 14: Percentage of smokers who have smoked in the past 12 months but report not smoking now. Base: All those who smoked in the past year and all those who smoked in the past year and made at least one quit attempt. $N=5,958(2007)$, $4,602(2008)$, $4,973(2009)$, $5,775(2010)$ for all smokers and $N=2,533(2007)$, $1,829(2008)$, $1,833(2009)$, $2,068(2009)$, $2,068(2010)$

Aggregating data over all years, a multiple logistic regression analysis was used to assess the predictors of current non-smoking status among those who had attempted to stop within the past 12 months. All predictors were entered simultaneously (giving adjusted odds ratios). Table 1 shows the results.

Variable	Odds ratio (adjusted)	95% confidence interval
Age (years)	1.15**	1.09-1.21
Routine and manual occupation vs non-routine and manual (reference)	0.76**	0.66-0.86
Male vs female (reference)	1.05	0.90-1.21
Dependence (1-6)	0.28**	0.26-0.31
Time since most recent quit attempt		
26-52 weeks (reference)	1	
<1 week	13.05**	9.56-17.82
1-4 weeks	3.92**	3.07-5.02
4-8 weeks	1.39*	1.07-1.81
8-12 weeks	1.16	0.88-1.51
12-26 weeks	1.17	0.95-1.44
Number of prior quit attempts in the past 12 months	0.64**	0.58-0.71
Abrupt vs gradual cessation (reference)	1.89**	1.60-2.22
Unplanned vs planned in advance (reference)	0.97	0.83-1.13
Method used		
Unaided (reference)	1	
NHS specialist support	3.53**	2.12-5.88
Medication Rx	1.73**	1.39-2.15
NRT OTC	0.97	0.81-1.16

Table 1: Predictor of success (currently not smoking) of quit attempts made in the previous 12 months by forced entry multiple logistic regression. Base=All respondents who had smoked in the past year and made at least one quit attempt. N=8,551. Routine and manual=social grade C-E; Dependence=ratings of strength of urges smoke (1=no urges-6=extremely strong); Abrupt=tried to quit without cutting down first; Unplanned=tried to quit as soon as made the decision; Rx=prescription with minimal behavioural support, NRT OTC=nicotine replacement therapy bought over the counter

Success was associated with greater age, non routine and manual occupation, lower dependence, not having tried to quit previously that year, having made the quit attempt more recently, and abrupt rather than gradual cessation. Use of specialist NHS support (which includes medication) was associated with 3.53 times the odds of quitting compared with unaided quit attempts and use of medication on prescription was associated with 1.72 times the odds of quitting. No improvement in success rates could be found for NRT bought over the counter.

A logistic regression analysis was undertaken restricted to respondents who had tried to quit with medication obtained on prescription or provided by the NHS specialist service to determine whether the type of medication used was associated with success rates. It was found that those who used varenicline had 1.48 (95% CI 1.03-2.14) times the odds of success compared with those using NRT after adjusting for all the predictor variables used previously.

References

1. Fidler JA, Shahab L, West O, et al. 'The Smoking Toolkit Study': A national study of smoking and smoking cessation in England. *BMC Public Health*. Jun 18 2011;11(1):479.
2. Beard E, Fidler J, West R. Is use of nicotine replacement therapy while continuing to smoke associated with increased nicotine intake? Evidence from a population sample. *Psychopharmacology (Berl)*. May 26 In Press.
3. Beard E, McNeill A, Aveyard P, Michie S, West R. Use of Nicotine Replacement Therapy for smoking reduction and during enforced temporary abstinence: a national survey of English smokers. *Addiction*. 2011;106(1):197-204.
4. Berg CJ, An LC, Kirch M, et al. Failure to report attempts to quit smoking. *Addict Behav*. Oct 2010;35(10):900-904.

5. Fidler J, Shahab L, West R. Strength of urges to smoke as a measure of severity of cigarette dependence: comparison with the Fagerstrom Test for Nicotine Dependence and its components. *Addiction*. 2011;106(3):631-638.
6. Fidler J, West R. Enjoyment of smoking and urges to smoke as predictors of attempts and success of attempts to stop smoking: a longitudinal study. *Drug and Alcohol Dependence* 2011;115(1-2):30-34.
7. Fidler JA, Stapleton JA, West R. Variation in saliva cotinine as a function of self-reported attempts to reduce cigarette consumption. *Psychopharmacology (Berl)*. May 10 2011.
8. Fidler JA, West R. Self-perceived smoking motives and their correlates in a general population sample. *Nicotine Tob Res*. Oct 2009;11(10):1182-1188.
9. Fidler JA, West R. Changes in smoking prevalence in 16-17-year-old versus older adults following a rise in legal age of sale: findings from an English population study. *Addiction*. Nov 2010;105(11):1984-1988.
10. Gardner B, West R. Public support in England for raising the price of cigarettes to fund tobacco control activities. *Tob Control*. Aug 2010;19(4):331-333.
11. Hackshaw L, McEwen A, West R, Bauld L. Quit attempts in response to smoke-free legislation in England. *Tob Control*. Apr 2010;19(2):160-164.
12. Kotz D, Fidler J, West R. Factors associated with the use of aids to cessation in English smokers. *Addiction*. Aug 2009;104(8):1403-1410.
13. Kotz D, Fidler JA, West R. Did the Introduction of Varenicline in England Substitute for or Add to the Use of Other Smoking Cessation Medications? *Nicotine Tob Res*. May 4 2011.
14. Kotz D, Stapleton J, Owen L, West R. How cost-effective is "No Smoking Day"? *Tob Control*. 2011;20(4):302-304. .
15. Kotz D, West R. Explaining the social gradient in smoking cessation: it's not in the trying, but in the succeeding. *Tob Control*. Feb 2009;18(1):43-46.
16. Shahab L, West R. Do ex-smokers report feeling happier following cessation? Evidence from a cross-sectional survey. *Nicotine Tob Res*. May 2009;11(5):553-557.
17. Shahab L, West R. Public support in England for a total ban on the sale of tobacco products. *Tob Control*. Apr 2010;19(2):143-147.
18. Smit ES, Fidler JA, West R. The role of desire, duty and intention in predicting attempts to quit smoking. *Addiction*. Dec 2 2011;106(4):844-851.
19. Vangeli E, West R. Sociodemographic differences in triggers to quit smoking: findings from a national survey. *Tob Control*. Dec 2008;17(6):410-415.